

**New nucleotide sequences which code for the menE gene**

The invention provides nucleotide sequences from coryneform bacteria which code for the menE gene and a process for the fermentative preparation of amino acids using bacteria in  
5 which the menE gene is attenuated.

**Prior art**

L-Amino acids, in particular L-lysine, are used in human medicine and in the pharmaceuticals industry, in the foodstuffs industry and very particularly in animal  
10 nutrition.

It is known that amino acids are prepared by fermentation from strains of coryneform bacteria, in particular *Corynebacterium glutamicum*. Because of their great importance, work is constantly being undertaken to improve  
15 the preparation processes. Improvements to the process can relate to fermentation measures, such as, for example, stirring and supply of oxygen, or the composition of the nutrient media, such as, for example, the sugar concentration during the fermentation, or the working up to  
20 the product form by, for example, ion exchange chromatography, or the intrinsic output properties of the microorganism itself.

Methods of mutagenesis, selection and mutant selection are used to improve the output properties of these  
25 microorganisms. Strains which are resistant to antimetabolites or are auxotrophic for metabolites of regulatory importance and which produce amino acids are obtained in this manner.

Methods of the recombinant DNA technique have also been  
30 employed for some years for improving the strain of *Corynebacterium* strains which produce L-amino acid, by amplifying individual amino acid biosynthesis genes and investigating the effect on the amino acid production.

## Object of the invention

The inventors had the object of providing new measures for improved fermentative preparation of amino acids.

## Description of the invention

- 5 Where L-amino acids or amino acids are mentioned in the following, this means one or more amino acids, including their salts, chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-  
10 isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine. L-Lysine is particularly preferred.

- When L-lysine or lysine are mentioned in the following, not only the bases but also the salts, such as e.g. lysine  
15 monohydrochloride or lysine sulfate, are meant by this.

The invention provides an isolated polynucleotide from coryneform bacteria, comprising a polynucleotide sequence which codes for the menE gene, chosen from the group consisting of

- 20 a) polynucleotide which is identical to the extent of at least 70% to a polynucleotide which codes for a polypeptide which comprises the amino acid sequence of SEQ ID No. 2,  
b) polynucleotide which codes for a polypeptide which  
25 comprises an amino acid sequence which is identical to the extent of at least 70% to the amino acid sequence of SEQ ID No. 2,  
c) polynucleotide which is complementary to the polynucleotides of a) or b), and

d) polynucleotide comprising at least 15 successive nucleotides of the polynucleotide sequence of a), b) or c),

the polypeptide preferably having the activity of O-succinylbenzoic acid CoA ligase.

The invention also provides the abovementioned polynucleotide, this preferably being a DNA which is capable of replication, comprising:

- (i) the nucleotide sequence, shown in SEQ ID No.1, or
- 10 (ii) at least one sequence which corresponds to sequence (i) within the range of the degeneration of the genetic code, or
- (iii) at least one sequence which hybridizes with the sequences complementary to sequences (i) or (ii),
- 15 and optionally
- (iv) sense mutations of neutral function in (i).

The invention also provides:

a polynucleotide, in particular DNA, which is capable of replication and comprises the nucleotide sequence as shown in SEQ ID No.1;

a polynucleotide which codes for a polypeptide which comprises the amino acid sequence as shown in SEQ ID No. 2;

a vector containing parts of the polynucleotide according to the invention, but at least 15 successive nucleotides of the sequence claimed,

and coryneform bacteria in which the menE gene is attenuated, in particular by an insertion or deletion.

The invention also provides polynucleotides which substantially comprise a polynucleotide sequence, which are obtainable by screening by means of hybridization of a corresponding gene library of a coryneform bacterium, which

5 comprises the complete gene or parts thereof, with a probe which comprises the sequence of the polynucleotide according to the invention according to SEQ ID No.1 or a fragment thereof, and isolation of the polynucleotide sequence mentioned.

- 10 Polynucleotides which comprise the sequences according to the invention are suitable as hybridization probes for RNA, cDNA and DNA, in order to isolate, in the full length, nucleic acids or polynucleotides or genes which code for O-succinylbenzoic acid CoA ligase or to isolate those nucleic
- 15 acids or polynucleotides or genes which have a high similarity with the sequence the menE gene.

- Polynucleotides which comprise the sequences according to the invention are furthermore suitable as primers with the aid of which DNA of genes which code for O-succinylbenzoic
- 20 acid CoA ligase can be prepared by the polymerase chain reaction (PCR).

- Such oligonucleotides which serve as probes or primers comprise at least 30, preferably at least 20, very particularly preferably at least 15 successive nucleotides.
- 25 Oligonucleotides which have a length of at least 40 or 50 nucleotides are also suitable.

"Isolated" means separated out of its natural environment.

- "Polynucleotide" in general relates to polyribonucleotides and polydeoxyribonucleotides, it being possible for these
- 30 to be non-modified RNA or DNA or modified RNA or DNA.

The polynucleotides according to the invention include a polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom and also those which are at least 70%,

preferably at least 80% and in particular at least 90% to 95% identical to the polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom.

- 5 "Polypeptides" are understood as meaning peptides or proteins which comprise two or more amino acids bonded via peptide bonds.

- 10 The polypeptides according to the invention include a polypeptide according to SEQ ID No. 2, in particular those with the biological activity of O-succinylbenzoic acid CoA ligase, and also those which are at least 70%, preferably at least 80% and in particular at least 90% to 95% identical to the polypeptide according to SEQ ID No. 2 and have the activity mentioned.

- 15 The invention furthermore relates to a process for the fermentative preparation of amino acids chosen from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine using coryneform bacteria which in particular  
20 already produce amino acids and in which the nucleotide sequences which code for the menE gene are attenuated, in particular eliminated or expressed at a low level.

- 25 The term "attenuation" in this connection describes the reduction or elimination of the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by using a weak promoter or using a gene or allele which codes for a corresponding enzyme with a low activity or inactivates the  
30 corresponding gene or enzyme (protein), and optionally combining these measures.

The microorganisms to which the present invention relates can prepare amino acids from glucose, sucrose, lactose,

fructose, maltose, molasses, starch, cellulose or from glycerol and ethanol. They can be representatives of coryneform bacteria, in particular of the genus *Corynebacterium*. Of the genus *Corynebacterium*, there may be mentioned in particular the species *Corynebacterium glutamicum*, which is known among experts for its ability to produce L-amino acids.

Suitable strains of the genus *Corynebacterium*, in particular of the species *Corynebacterium glutamicum* (*C. glutamicum*), are in particular the known wild-type strains

*Corynebacterium glutamicum* ATCC13032  
*Corynebacterium acetoglutamicum* ATCC15806  
*Corynebacterium acetoacidophilum* ATCC13870  
*Corynebacterium melassecola* ATCC17965  
*Corynebacterium thermoaminogenes* FERM BP-1539  
*Brevibacterium flavum* ATCC14067  
*Brevibacterium lactofermentum* ATCC13869 and  
*Brevibacterium divaricatum* ATCC14020

and L-amino acid-producing mutants or strains prepared therefrom.

The new *menE* gene from *C. glutamicum* which codes for the enzyme O-succinylbenzoic acid CoA ligase (EC 6.2.1.26) has been isolated.

To isolate the *menE* gene or also other genes of *C.*

*glutamicum*, a gene library of this microorganism is first set up in *Escherichia coli* (*E. coli*). The setting up of gene libraries is described in generally known textbooks and handbooks. The textbook by Winnacker: *Gene und Klone, Eine Einführung in die Gentechnologie* (Verlag Chemie, Weinheim, Germany, 1990), or the handbook by Sambrook et al.: *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989) may be mentioned as an example. A well-known gene library is that of the *E. coli*

K-12 strain W3110 set up in  $\lambda$  vectors by Kohara et al. (Cell 50, 495-508 (1987)). Bathe et al. (Molecular and General Genetics, 252:255-265, 1996) describe a gene library of *C. glutamicum* ATCC13032, which was set up with the aid of the cosmid vector SuperCos I (Wahl et al., 1987, Proceedings of the National Academy of Sciences USA, 84:2160-2164) in the *E. coli* K-12 strain NM554 (Raleigh et al., 1988, Nucleic Acids Research 16:1563-1575).

- 5  
10 B6rmann et al. (Molecular Microbiology 6(3), 317-326) (1992)) in turn describe a gene library of *C. glutamicum* ATCC13032 using the cosmid pHc79 (Hohn and Collins, 1980, Gene 11, 291-298).

To prepare a gene library of *C. glutamicum* in *E. coli* it is also possible to use plasmids such as pBR322 (Bolivar, 1979, Life Sciences, 25, 807-818) or pUC9 (Vieira et al., 1982, Gene, 19:259-268). Suitable hosts are, in particular, those *E. coli* strains which are restriction- and recombination-defective, such as, for example, the strain DH5 $\alpha$ mc<sub>r</sub>, which has been described by Grant et al.

- 15  
20 (Proceedings of the National Academy of Sciences USA, 87 (1990) 4645-4649). The long DNA fragments cloned with the aid of cosmids or other  $\lambda$  vectors can then in turn be subcloned and subsequently sequenced in the usual vectors which are suitable for DNA sequencing, such as is described  
25 e. g. by Sanger et al. (Proceedings of the National Academy of Sciences of the United States of America, 74:5463-5467, 1977).

- The resulting DNA sequences can then be investigated with known algorithms or sequence analysis programs, such as  
30 e.g. that of Staden (Nucleic Acids Research 14, 217-232(1986)), that of Marck (Nucleic Acids Research 16, 1829-1836 (1988)) or the GCG program of Butler (Methods of Biochemical Analysis 39, 74-97 (1998)).

00034752.02301

5

10

15

20

25

30

35



Instructions for identifying DNA sequences by means of hybridization can be found by the expert, inter alia, in the handbook "The DIG System Users Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim, Germany, 1993) and in Liebl et al. (International Journal of Systematic Bacteriology 41: 255-260 (1991)). The hybridization takes place under stringent conditions, that is to say only hybrids in which the probe and target sequence, i. e. the polynucleotides treated with the probe, are at least 70% identical are formed. It is known that the stringency of the hybridization, including the washing steps, is influenced or determined by varying the buffer composition, the temperature and the salt concentration. The hybridization reaction is preferably carried out under a relatively low stringency compared with the washing steps (Hybaid Hybridisation Guide, Hybaid Limited, Teddington, UK, 1996).

A 5x SSC buffer at a temperature of approx. 50°C - 68°C, for example, can be employed for the hybridization reaction. Probes can also hybridize here with polynucleotides which are less than 70% identical to the sequence of the probe. Such hybrids are less stable and are removed by washing under stringent conditions. This can be achieved, for example, by lowering the salt concentration to 2x SSC and optionally subsequently 0.5x SSC (The DIG System User's Guide for Filter Hybridisation, Boehringer Mannheim, Mannheim, Germany, 1995) a temperature of approx. 50°C - 68°C being established. It is optionally possible to lower the salt concentration to 0.1x SSC. Polynucleotide fragments which are, for example, at least 70% or at least 80% or at least 90% to 95% identical to the sequence of the probe employed can be isolated by increasing the hybridization temperature stepwise from 50°C to 68°C in steps of approx. 1 - 2°C. Further instructions on hybridization are obtainable on the market in the form of

05834725-072301

so-called kits (e.g. DIG Easy Hyb from Roche Diagnostics GmbH, Mannheim, Germany, Catalogue No. 1603558).

Instructions for amplification of DNA sequences with the aid of the polymerase chain reaction (PCR) can be found by the expert, inter alia, in the handbook by Gait:  
5 Oligonukleotide [sic] synthesis: A Practical Approach (IRL Press, Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994).

It has been found that coryneform bacteria produce amino acids in an improved manner after attenuation of the menE gene.

To achieve an attenuation, either the expression of the menE gene or the catalytic properties of the enzyme protein can be reduced or eliminated. The two measures can  
15 optionally be combined.

The reduction in gene expression can take place by suitable culturing or by genetic modification (mutation) of the signal structures of gene expression. Signal structures of gene expression are, for example, repressor genes,  
20 activator genes, operators, promoters, attenuators, ribosome binding sites, the start codon and terminators. The expert can find information on this e.g. in the patent application WO 96/15246, in Boyd and Murphy (Journal of Bacteriology 170: 5949 (1988)), in Voskuil and Chambliss  
25 (Nucleic Acids Research 26: 3548 (1998)), in Jensen and Hammer (Biotechnology and Bioengineering 58: 191 (1998)), in Pátek et al. (Microbiology 142: 1297 (1996)), Vasicova et al. (Journal of Bacteriology 181: 6188 (1999)) and in known textbooks of genetics and molecular biology, such as  
30 e.g. the textbook by Knippers ("Molekulare Genetik", 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995) or that by Winnacker ("Gene und Klone ", VCH Verlagsgesellschaft, Weinheim, Germany, 1990).

Mutations which lead to a change or reduction in the catalytic properties of enzyme proteins are known from the prior art; examples which may be mentioned are the works by Qiu and Goodman (Journal of Biological Chemistry 272: 8611-8617 (1997)), Sugimoto et al. (Bioscience Biotechnology and Biochemistry 61: 1760-1762 (1997)) and Möckel ("Die Threonindehydratase aus Corynebacterium glutamicum: Aufhebung der allosterischen Regulation und Struktur des Enzyms", Reports from the Jülich Research Centre, Jül-2906, ISSN09442952, Jülich, Germany, 1994). Summarizing descriptions can be found in known textbooks of genetics and molecular biology, such as e.g. that by Hagemann ("Allgemeine Genetik", Gustav Fischer Verlag, Stuttgart, 1986).

Possible mutations are transitions, transversions, insertions and deletions. Depending on the effect of the amino acid exchange on the enzyme activity, "missense mutations" or "nonsense mutations" are referred to. Insertions or deletions of at least one base pair (bp) in a gene lead to frame shift mutations, as a consequence of which incorrect amino acids are incorporated or translation is interrupted prematurely. Deletions of several codons typically lead to a complete loss of the enzyme activity. Instructions on generation of such mutations are prior art and can be found in known textbooks of genetics and molecular biology, such as e.g. the textbook by Knippers ("Molekulare Genetik", 6th edition, Georg Thieme Verlag, Stuttgart, Germany, 1995), that by Winnacker ("Gene und Klone", VCH Verlagsgesellschaft, Weinheim, Germany, 1990) or that by Hagemann ("Allgemeine Genetik", Gustav Fischer Verlag, Stuttgart, 1986).

A common method of mutating genes of *C. glutamicum* is the method of "gene disruption" and "gene replacement" described by Schwarzer and Pühler (Bio/Technology 9, 84-87 (1991)).

In the method of gene disruption a central part of the coding region of the gene of interest is cloned in a plasmid vector which can replicate in a host (typically *E. coli*), but not in *C. glutamicum*. Possible vectors are, for example, pSUP301 (Simon et al., *Bio/Technology* 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., *Gene* 145, 69-73 (1994)), pK18mobsacB or pK19mobsacB (Jäger et al., *Journal of Bacteriology* 174: 5462-65 (1992)), pGEM-T (Promega corporation, Madison, WI, USA), pCR2.1-TOPO (Shuman (1994). *Journal of Biological Chemistry* 269:32678-84; US Patent 5,487,993), pCR@Blunt (Invitrogen, Groningen, Holland; Bernard et al., *Journal of Molecular Biology*, 234: 534-541 (1993)) or pEM1 (Schrumpf et al, 1991, *Journal of Bacteriology* 173:4510-4516). The plasmid vector which contains the central part of the coding region of the gene is then transferred into the desired strain of *C. glutamicum* by conjugation or transformation. The method of conjugation is described, for example, by Schäfer et al. (*Applied and Environmental Microbiology* 60, 756-759 (1994)). Methods for transformation are described, for example, by Thierbach et al. (*Applied Microbiology and Biotechnology* 29, 356-362 (1988)), Dunican and Shivanan (*Bio/Technology* 7, 1067-1070 (1989)) and Tauch et al. (*FEMS Microbiological Letters* 123, 343-347 (1994)). After homologous recombination by means of a "cross-over" event, the coding region of the gene in question is interrupted by the vector sequence and two incomplete alleles are obtained, one lacking the 3' end and one lacking the 5' end. This method has been used, for example, by Fitzpatrick et al. (*Applied Microbiology and Biotechnology* 42, 575-580 (1994)) to eliminate the *recA* gene of *C. glutamicum*.

In the method of "gene replacement", a mutation, such as e.g. a deletion, insertion or base exchange, is established in vitro in the gene of interest. The allele prepared in its turn cloned in a vector which is not replicative for *C. glutamicum* and this is then transferred into the desired

host of *C. glutamicum* by transformation or conjugation. After homologous recombination by means of a first "cross-over" event which effects integration and a suitable second "cross-over" event which effects excision in the target gene or in the target sequence, the incorporation of the mutation or of the allele is achieved. This method was used, for example, by Peters-Wendisch et al. (Microbiology 144, 915 - 927 (1998)) to eliminate the *pyc* gene of *C. glutamicum* by a deletion.

- 5
- 10 A deletion, insertion or a base exchange can be incorporated into the *menE* gene in this manner.

In addition, it may be advantageous for the production of L-amino acids to enhance, in particular over-express, one or more enzymes of the particular biosynthesis pathway, of glycolysis, of anaplerosis, of the citric acid cycle, of the pentose phosphate cycle, of amino acid export and optionally regulatory proteins, in addition to the attenuation of the *menE* gene.

- 15
- 20 The term "enhancement" in this connection describes the increase in the intracellular activity of one or more enzymes (proteins) in a microorganism which are coded by the corresponding DNA, for example by increasing the number of copies of the gene or genes, using a potent promoter or using a gene or allele which codes for a corresponding
- 25 enzyme (protein) having a high activity, and optionally combining these measures.

Thus, for the preparation of L-amino acids, in addition to the attenuation of the *menE* gene at the same time one or more of the genes chosen from the group consisting of

- 30 • the *dapA* gene which codes for dihydrodipicolinate synthase (EP-B 0 197 335),

- the gap gene which codes for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the tpi gene which codes for triose phosphate isomerase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the pgk gene which codes for 3-phosphoglycerate kinase (Eikmanns (1992), Journal of Bacteriology 174:6076-6086),
- the zwf gene which codes for glucose 6-phosphate dehydrogenase (JP-A-09224661),
- 10 • the pyc gene which codes for pyruvate carboxylase (DE-A-198 31 609),
- the mqo gene which codes for malate-quinone oxidoreductase (Molenaar et al., European Journal of Biochemistry 254, 395-403 (1998)),
- 15 • the lysC gene which codes for a feed-back resistant aspartate kinase (Accession No.P26512; EP-B-0387527; EP-A-0699759; WO 00/63388),
- the lysE gene which codes for lysine export (DE-A-195 48 222),
- 20 • the hom gene which codes for homoserine dehydrogenase (EP-A 0131171),
- the ilvA gene which codes for threonine dehydratase (Möckel et al., Journal of Bacteriology (1992) 8065-8072) or the ilvA(Fbr) allele which codes for a "feed back resistant" threonine dehydratase (Möckel et al.,
- 25 (1994) Molecular Microbiology 13: 833-842),
- the ilvBN gene which codes for acetohydroxy-acid synthase (EP-B 0356739),

- the *ilvD* gene which codes for dihydroxy-acid dehydratase (Sahm and Eggeling (1999) Applied and Environmental Microbiology 65: 1973-1979),

- the *zwa1* gene which codes for the Zwa1 protein (DE: 19959328.0, DSM 13115)

can be enhanced, in particular over-expressed.

It may furthermore be advantageous for the production of amino acids, in addition to the attenuation of the *menE* gene, at the same time for one or more of the genes chosen from the group consisting of

- the *pck* gene which codes for phosphoenol pyruvate carboxykinase (DE 199 50 409.1, DSM 13047),
- the *pgi* gene which codes for glucose 6-phosphate isomerase (US 09/396,478, DSM 12969),
- the *poxB* gene which codes for pyruvate oxidase (DE:1995 1975.7, DSM 13114),
- the *zwa2* gene which codes for the Zwa2 protein (DE: 19959327.2, DSM 13113)

to be attenuated, in particular for the expression thereof to be reduced.

In addition to the attenuation of the *menE* gene it may furthermore be advantageous for the production of amino acids to eliminate undesirable side reactions (Nakayama: "Breeding of Amino Acid Producing Microorganisms", in: Overproduction of Microbial Products, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The invention also provides the microorganisms prepared according to the invention, and these can be cultured continuously or discontinuously in the batch process (batch culture) or in the fed batch (feed process) or repeated fed

batch process (repetitive feed process) for the purpose of production of L-amino acids. A summary of known culture methods is described in the textbook by Chmiel

(Bioprozesstechnik 1. Einführung in die

- 5 Bioverfahrenstechnik (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen (Vieweg Verlag, Braunschweig/Wiesbaden, 1994)).

The culture medium to be used must meet the requirements of

10 the particular strains in a suitable manner. Descriptions of culture media for various microorganisms are contained in the handbook "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

- 15 Sugars and carbohydrates, such as e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, such as, for example, soya oil, sunflower oil, groundnut oil and coconut fat, fatty acids, such as, for example, palmitic acid, stearic acid and linoleic acid,
- 20 alcohols, such as, for example, glycerol and ethanol, and organic acids, such as, for example, acetic acid, can be used as the source of carbon. These substances can be used individually or as a mixture.

- Organic nitrogen-containing compounds, such as peptones,
- 25 yeast extract, meat extract, malt extract, corn steep liquor, soya bean flour and urea, or inorganic compounds, such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate, can be used as the source of nitrogen. The sources of nitrogen can
- 30 be used individually or as a mixture.

Phosphoric acid, potassium dihydrogen phosphate or dipotassium hydrogen phosphate or the corresponding sodium-containing salts can be used as the source of phosphorus. The culture medium must furthermore comprise salts of



metals, such as, for example, magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth substances, such as amino acids and vitamins, can be employed in addition to the abovementioned substances.

- 5 Suitable precursors can moreover be added to the culture medium. The starting substances mentioned can be added to the culture in the form of a single batch, or can be fed in during the culture in a suitable manner.

- Basic compounds, such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acid compounds, such as phosphoric acid or sulfuric acid, can be employed in a suitable manner to control the pH of the culture. Antifoams, such as, for example, fatty acid polyglycol esters, can be employed to control the development of foam.
- 10 Suitable substances having a selective action, such as, for example, antibiotics, can be added to the medium to maintain the stability of plasmids. To maintain aerobic conditions, oxygen or oxygen-containing gas mixtures, such as, for example, air, are introduced into the culture. The
- 15 temperature of the culture is usually 20°C to 45°C, and preferably 25°C to 40°C. Culturing is continued until a maximum of the desired product has formed. This target is usually reached within 10 hours to 160 hours.
- 20

- Methods for the determination of L-amino acids are known from the prior art. The analysis can thus be carried out, for example, as described by Spackman et al. (Analytical Chemistry, 30, (1958), 1190) by anion exchange chromatography with subsequent ninhydrin derivatization, or it can be carried out by reversed phase HPLC, for example
- 25 as described by Lindroth et al. (Analytical Chemistry (1979) 51: 1167-1174).
- 30

The process according to the invention is used for fermentative preparation of amino acids.

The following microorganism was deposited on 26.02.2001 as a pure culture at the Deutsche Sammlung für Mikroorganismen [sic] und Zellkulturen (DSMZ = German Collection of Microorganisms and Cell Cultures, Braunschweig, Germany) in accordance with the Budapest Treaty:

- Escherichia coli Top10/pCR2.1menEint as DSM 14080.

The present invention is explained in more detail in the following with the aid of embodiment examples.

- 10 The isolation of plasmid DNA from Escherichia coli and all techniques of restriction, Klenow and alkaline phosphatase treatment were carried out by the method of Sambrook et al. (Molecular Cloning. A Laboratory Manual, 1989, Cold Spring Harbour Laboratory Press, Cold Spring Harbor, NY, USA).
- 15 Methods for transformation of Escherichia coli are also described in this handbook.

The composition of the usual nutrient media, such as LB or TY medium, can also be found in the handbook by Sambrook et al.

#### Example 1

- 20 Preparation of a genomic cosmid gene library from C. glutamicum ATCC 13032

- Chromosomal DNA from C. glutamicum ATCC 13032 was isolated as described by Tauch et al. (1995, Plasmid 33:168-179) and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Code no. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Code no. 1758250). The DNA of the cosmid vector SuperCos1 (Wahl et al. (1987), Proceedings of the National Academy of Sciences, USA 84:2160-2164), obtained from Stratagene (La Jolla, USA, Product Description
- 25
- 30

SuperCosI Cosmid Vector Kit, Code no. 251301) was cleaved with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, Product Description XbaI, Code no. 27-0948-02) and likewise dephosphorylated with shrimp alkaline phosphatase.

The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Code no. 27-0868-04). The cosmid DNA treated in this manner was mixed with the treated ATCC13032 DNA and the batch was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, Product Description T4-DNA-Ligase, Code no.27-0870-04). The ligation mixture was then packed in phages with the aid of Gigapack II XL Packing Extract (Stratagene, La Jolla, USA, Product Description Gigapack II XL Packing Extract, Code no. 200217).

For infection of the E. coli strain NM554 (Raleigh et al. 1988, Nucleic Acid Res. 16:1563-1575) the cells were taken up in 10 mM MgSO<sub>4</sub> and mixed with an aliquot of the phage suspension. The infection and titering of the cosmid library were carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the cells being plated out on LB agar (Lennox, 1955, Virology, 1:190) + 100 µg/ml ampicillin. After incubation overnight at 37°C, recombinant individual clones were selected.

### Example 2

#### Isolation and sequencing of the menE gene

The cosmid DNA of an individual colony was isolated with the Qiaprep Spin Miniprep Kit (Product No. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partly cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, Product Description Sau3AI, Product No. 27-0913-02). The DNA

fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Molecular Biochemicals, Mannheim, Germany, Product Description SAP, Product No. 1758250). After separation by gel electrophoresis, the cosmid fragments in the size range of 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (Product No. 20021, Qiagen, Hilden, Germany).

The DNA of the sequencing vector pZero-1, obtained from Invitrogen (Groningen, The Netherlands, Product Description Zero Background Cloning Kit, Product No. K2500-01) was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, Product Description BamHI, Product No. 27-0868-04). The ligation of the cosmid fragments in the sequencing vector pZero-1 was carried out as described by Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then electroporated (Tauch et al. 1994, FEMS Microbiol. Letters, 123:343-7) into the E. coli strain DH5 $\alpha$ mc<sup>r</sup> (Grant, 1990, Proceedings of the National Academy of Sciences, U.S.A., 87:4645-4649). Letters, 123:343-7) and plated out on LB agar (Lennox, 1955, Virology, 1:190) with 50  $\mu$ g/ml zeocin.

The plasmid preparation of the recombinant clones was carried out with Biorobot 9600 (Product No. 900200, Qiagen, Hilden, Germany). The sequencing was carried out by the dideoxy chain termination method of Sanger et al. (1977, Proceedings of the National Academies of Sciences, U.S.A., 74:5463-5467) with modifications according to Zimmermann et al. (1990, Nucleic Acids Research, 18:1067). The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (Product No. 403044, Weiterstadt, Germany) was used. The separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphoresis NF Acrylamide/Bisacrylamide" Gel (29:1) (Product No.

A124.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).

- 5 The raw sequence data obtained were then processed using the Staden program package (1986, Nucleic Acids Research, 14:217-231) version 97-0. The individual sequences of the pZerol derivatives were assembled to a continuous contig. The computer-assisted coding region analysis [sic] were prepared with the XNIP program (Staden, 1986, Nucleic Acids Research, 14:217-231). Further analyses were carried out with the "BLAST search program" (Altschul et al., 1997, Nucleic Acids Research, 25:3389-3402) against the non-redundant databank of the "National Center for Biotechnology Information" (NCBI, Bethesda, MD, USA).
- 10
- 15 The resulting nucleotide sequence is shown in SEQ ID No. 1. Analysis of the nucleotide sequence showed an open reading frame of 1131 bp, which was called the menE gene. The menE gene codes for a polypeptide of 376 amino acids.

### Example 3

- 20 Preparation of an integration vector for integration mutagenesis of the menE gene

From the strain ATCC 13032, chromosomal DNA was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 - 1828 (1994)). On the basis of the sequence of the menE gene known for *C. glutamicum* from example 2, the following oligonucleotides were chosen for the polymerase chain reaction (see SEQ ID No. 3 and SEQ ID No. 4):

25

menE-int1:

5' CTC ACT CCG TTG AAT TTG G 3'

30 menE-int2:

5' CAG GTG CAT TTC TGT AGC C 3'

- The primers shown were synthesized by MWG Biotech (Ebersberg, Germany) and the PCR reaction was carried out by the standard PCR method of Innis et al. (PCR protocols. A guide to methods and applications, 1990, Academic Press)
- 5 with the Taq-polymerase from Boehringer Mannheim (Germany, Product Description Taq DNA polymerase, Product No. 1 146 165). With the aid of the polymerase chain reaction, the primers allow amplification of an internal fragment of the menE gene 520 bp in size. The product amplified in this way
- 10 was tested electrophoretically in a 0.8% agarose gel.

The amplified DNA fragment was ligated with the TOPO TA Cloning Kit from Invitrogen Corporation (Carlsbad, CA, USA; Catalogue Number K4500-01) in the vector pCR2.1-TOPO (Mead at al. (1991) Bio/Technology 9:657-663).

- 15 The E. coli strain TOP10 was then electroporated with the ligation batch (Hanahan, In: DNA cloning. A practical approach. Vol.I. IRL-Press, Oxford, Washington DC, USA, 1985). Selection of plasmid-carrying cells was carried out by plating out the transformation batch on LB Agar
- 20 (Sambrook et al., Molecular cloning: a laboratory manual. 2<sup>nd</sup> Ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989), which had been supplemented with 50 mg/l kanamycin. Plasmid DNA was isolated from a transformant with the aid of the QIAprep Spin Miniprep Kit
- 25 from Qiagen and checked by restriction with the restriction enzyme EcoRI and subsequent agarose gel electrophoresis (0.8%). The plasmid was called pCR2.1menEint and is shown in figure 1.

#### Example 4

- 30 Integration mutagenesis of the menE gene in the strain DSM 5715

The vector pCR2.1menEint mentioned in example 3 was electroporated by the electroporation method of Tauch et

al. (FEMS Microbiological Letters, 123:343-347 (1994)) in Corynebacterium glutamicum DSM 5715. The strain DSM 5715 is an AEC-resistant lysine producer. The vector pCR2.1menEint cannot replicate independently in DSM5715 and is retained  
5 in the cell only if it has integrated into the chromosome of DSM 5715. Selection of clones with pCR2.1menEint integrated into the chromosome was carried out by plating out the electroporation batch on LB agar (Sambrook et al., Molecular cloning: a laboratory manual. 2<sup>nd</sup> Ed. Cold Spring  
10 Harbor Laboratory Press, Cold Spring Harbor, N.Y.), which had been supplemented with 15 mg/l kanamycin.

For detection of the integration, the menEint fragment was labelled with the Dig hybridization kit from Boehringer by the method of "The DIG System Users Guide for Filter  
15 Hybridization" of Boehringer Mannheim GmbH (Mannheim, Germany, 1993). Chromosomal DNA of a potential integrant was isolated by the method of Eikmanns et al. (Microbiology 140: 1817 - 1828 (1994)) and in each case cleaved with the restriction enzymes EcoRI and PstI. The fragments formed  
20 were separated by means of agarose gel electrophoresis and hybridized at 68°C with the Dig hybridization [sic] kit from Boehringer. The plasmid pCR2.1menEint mentioned in example 3 had been inserted into the chromosome of DSM5715 within the chromosomal menE gene. The strain was called  
25 DSM5715::pCR2.1menEint.

#### Example 5

##### Preparation of lysine

The C. glutamicum strain DSM5715::pCR2.1menEint obtained in example 4 was cultured in a nutrient medium suitable for  
30 the production of lysine and the lysine content in the culture supernatant was determined.

For this, the strain was first incubated on an agar plate with the corresponding antibiotic (brain-heart agar with

kanamycin (25 mg/l) for 24 hours at 33°C. Starting from this agar plate culture, a preculture was seeded (10 ml medium in a 100 ml conical flask). The complete medium CgIII was used as the medium for the preculture.

2025-07-22 14:39:09





Medium Cg III

NaCl 2.5 g/l

Bacto-Peptone 10 g/l

Bacto-Yeast extract 10 g/l

Glucose (autoclaved separately) 2% (w/v)

The pH was brought to pH 7.4

5 Kanamycin (25 mg/l) was added to this. The preculture was incubated for 16 hours at 33°C at 240 rpm on a shaking machine. A main culture was seeded from this preculture such that the initial OD (660 nm) of the main culture was 0.1 OD. Medium MM was used for the main culture.

## Medium MM

CSL (corn steep liquor)	5 g/l
MOPS (morpholinopropanesulfonic acid)	20 g/l
Glucose (autoclaved separately)	50g/l
Salts:	
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub> )	25 g/l
KH <sub>2</sub> PO <sub>4</sub>	0.1 g/l
MgSO <sub>4</sub> * 7 H <sub>2</sub> O	1.0 g/l
CaCl <sub>2</sub> * 2 H <sub>2</sub> O	10 mg/l
FeSO <sub>4</sub> * 7 H <sub>2</sub> O	10 mg/l
MnSO <sub>4</sub> * H <sub>2</sub> O	5.0mg/l
Biotin (sterile-filtered)	0.3 mg/l
Thiamine * HCl (sterile-filtered)	0.2 mg/l
Leucine (sterile-filtered)	0.1 g/l
CaCO <sub>3</sub>	25 g/l

5 The CSL, MOPS and the salt solution are brought to pH 7 with aqueous ammonia and autoclaved. The sterile substrate and vitamin solutions are then added, and the CaCO<sub>3</sub> autoclaved in the dry state is added.

10 Culturing is carried out in a 10 ml volume in a 100 ml conical flask with baffles. Kanamycin (25 mg/l) was added. Culturing was carried out at 33°C and 80% atmospheric humidity.

- After 72 hours, the OD was determined at a measurement wavelength of 660 nm with a Biomek 1000 (Beckmann Instruments GmbH, Munich). The amount of lysine formed was determined with an amino acid analyzer from Eppendorf-
- 5 BioTronik (Hamburg, Germany) by ion exchange chromatography and post-column derivatization with ninhydrin detection.

The result of the experiment is shown in table 1.

Table 1

Strain	OD(660)	Lysine HCl g/l
DSM5715	8.2	12.64
DSM5715::pCR2.1menEint	9.2	14.89

# 10 Description of the figure:

Figure 1: Map of the plasmid pCR2.1menEint.

The abbreviations and designations used have the following meaning.

- KmR: Kanamycin resistance gene
- EcoRI: Cleavage site of the restriction enzyme EcoRI
- PstI: Cleavage site of the restriction enzyme PstI
- menEint: Internal fragment of the menE gene
- ColE1: Replication origin of the plasmid ColE1

## SEQUENCE PROTOCOL

&lt;110&gt; Degussa AG

5 &lt;120&gt; New nucleotide sequences which code for the menE gene

&lt;130&gt; 000551 BT

&lt;140&gt;

10 &lt;141&gt;

&lt;160&gt; 4

&lt;170&gt; PatentIn Ver. 2.1

15

&lt;210&gt; 1

&lt;211&gt; 1570

&lt;212&gt; DNA

&lt;213&gt; Corynebacterium glutamicum

20

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (230)..(1357)

&lt;223&gt; menE gene

25

&lt;400&gt; 1

ttcgttgcca tagacatgct cttcgagcga ctgtttgcgc acgtctcttc cggcatcttt 60

gtcaccaca atggttggga actcaccggc gcaatcgcg ctggcgcgct gcttctcacc 120

30

gcagttggcg caggtgcatg gagcatcgac ggggttctgg caaacgcaa ggccataaat 180

tagcgccaca actccgaatt ctgaaccatc ggcactagaa tctcggaat atg aat act 238

35

Met Asn Thr

1

cgc gtc ctc gaa gca cta cct gtt gat ctt gca gat ccc acc gca att 286

Arg Val Leu Glu Ala Leu Pro Val Asp Leu Ala Asp Pro Thr Ala Ile

40

5

10

15

ctg gga gat ctc gag gac gca atc tct ggg aag aaa act ttc ctc ccc 334

Leu Gly Asp Leu Glu Asp Ala Ile Ser Gly Lys Lys Thr Phe Leu Pro

50

20

25

30

35

atc cct gta caa gat aaa acc cgt gca cag ttg ctg cgc gat tct caa 382

Ile Pro Val Gln Asp Lys Thr Arg Ala Gln Leu Leu Arg Asp Ser Gln

45

40

45

50

cga gtt ggc ctc gcc atc gat cct tcg atc gct ttg gtg atg gcc act 430

50

Arg Val Gly Leu Ala Ile Asp Pro Ser Ile Ala Leu Val Met Ala Thr

55

60

65

tct ggt tct aca ggt acc ccg aag ggc gct cag ctc act ccg ttg aat 478

55

Ser Gly Ser Thr Gly Thr

70

75

80

ttg gtg agt tcc gcc gat gct acg cat cag ttt tta ggt ggc gaa ggc 526  
 Leu Val Ser Ser Ala Asp Ala Thr His Gln Phe Leu Gly Gly Glu Gly  
 85 90 95

5 cag tgg ttg ctt gcc atg cca gca cac cac att gca ggc atg cag gtg 574  
 Gln Trp Leu Leu Ala Met Pro Ala His His Ile Ala Gly Met Gln Val  
 100 105 110 115

10 ctt ctt cga agc ctc att gct gga gtt gag cca cta gct att gat ctc 622  
 Leu Leu Arg Ser Leu Ile Ala Gly Val Glu Pro Leu Ala Ile Asp Leu  
 120 125 130

15 agc aca ggt ttt cac att gac gct ttc gca ggc gcc gcg gca gaa ctg 670  
 Ser Thr Gly Phe His Ile Asp Ala Phe Ala Gly Ala Ala Ala Glu Leu  
 135 140 145

20 aaa aat acc ggc gac cgc gtc tat aca tcc ttg act cca atg cag tta 718  
 Lys Asn Thr Gly Asp Arg Val Tyr Thr Ser Leu Thr Pro Met Gln Leu  
 150 155 160

ctt aaa gca atg gac tcc ttg caa ggc att gaa gcc ctg aaa ctt ttt 766  
 Leu Lys Ala Met Asp Ser Leu Gln Gly Ile Glu Ala Leu Lys Leu Phe  
 165 170 175

25 gat gtc att ctt gtt ggc ggt gct gca ttg tct aag cag gcc cga att 814  
 Asp Val Ile Leu Val Gly Gly Ala Ala Leu Ser Lys Gln Ala Arg Ile  
 180 185 190 195

30 tct gcg gag cag cta gac atc aac att gtc acc acc tac ggc tcc tca 862  
 Ser Ala Glu Gln Leu Asp Ile Asn Ile Val Thr Thr Tyr Gly Ser Ser  
 200 205 210

35 gag act tca ggt ggc tgc gtt tat gat ggc aag ccc att ccc ggc gcg 910  
 Glu Thr Ser Gly Gly Cys Val Tyr Asp Gly Lys Pro Ile Pro Gly Ala  
 215 220 225

40 aaa gtc cgt att tcg gat gag cgc att gag ttg ggt ggc ccg atg att 958  
 Lys Val Arg Ile Ser Asp Glu Arg Ile Glu Leu Gly Gly Pro Met Ile  
 230 235 240

gcg cag ggc tac aga aat gca cct gaa cat ccg gat ttc gcc aac gag 1006  
 Ala Gln Gly Tyr Arg Asn Ala Pro Glu His Pro Asp Phe Ala Asn Glu  
 245 250 255

45 ggt tgg ttt acc acc tct gat tca ggt gaa ctc cac gac ggg att ctc 1054  
 Gly Trp Phe Thr Thr Ser Asp Ser Gly Glu Leu His Asp Gly Ile Leu  
 260 265 270 275

50 acc gtg act ggt cgc gtg gat acc gtc att gat tcc ggt gga ttg aag 1102  
 Thr Val Thr Gly Arg Val Asp Thr Val Ile Asp Ser Gly Gly Leu Lys  
 280 285 290

55 ttg cac cca gag gta ctg gaa cgt gcc atc gca gat att aaa ggt gtc 1150  
 Leu His Pro Glu Val Leu Glu Arg Ala Ile Ala Asp Ile Lys Gly Val  
 295 300 305

acc gcg gcg tgt gtt gtg ggt att ccc gat ccc cga tta ggc caa gca 1198  
 Thr Ala Ala Cys Val Val Gly Ile Pro Asp Pro Arg Leu Gly Gln Ala  
 310 315 320

att gtg gcc gcg tac tcc gga tgg atc agt ccg tct gaa gtt att gaa 1246  
 Ile Val Ala Ala Tyr Ser Gly Ser Ile Ser Pro Ser Glu Val Ile Glu  
 325 330 335  
 5 ggc ctc gac gat cta cct cgt tgg cag ctt ccc aaa cgg ctg aag cat 1294  
 Gly Leu Asp Asp Leu Pro Arg Trp Gln Leu Pro Lys Arg Leu Lys His  
 340 345 350 355  
 10 ctg gaa tct ttg ccc agc att ggt cct gga aaa gct gat cga cgt gct 1342  
 Leu Glu Ser Leu Pro Ser Ile Gly Pro Gly Lys Ala Asp Arg Arg Ala  
 360 365 370  
 15 atc gcg aag ctg ttt tagtcttcat tcttctgtggc tgcaactagt tttgccacat 1397  
 Ile Ala Lys Leu Phe  
 375  
 cttcatcggt gtacactttg gcgatctgct catcatttcc acccatgagg gtgttgccaa 1457  
 20 caactagtgc tcccaacttg gtggtgggca cgacagcgaa gtgtcggggc tgagcgtaga 1517  
 cctggcgaat aggggtgatca gagcgacgtg cgcaggcatg cagccatacg tca 1570  
 25 <210> 2  
 <211> 376  
 <212> PRT  
 <213> Corynebacterium glutamicum  
 30 <400> 2  
 Met Asn Thr Arg Val Leu Glu Ala Leu Pro Val Asp Leu Ala Asp Pro  
 1 5 10 15  
 35 Thr Ala Ile Leu Gly Asp Leu Glu Asp Ala Ile Ser Gly Lys Lys Thr  
 20 25  
 Phe Leu Pro Ile Pro Val Gln Asp Lys Thr Arg Ala Gln Leu Leu Arg  
 35 40 45  
 40 Asp Ser Gln Arg Val Gly Leu Ala Ile Asp Pro Ser Ile Ala Leu Val  
 50 55 60  
 Met Ala Thr Ser Gly Ser Thr Gly Thr Pro Lys Gly Ala Gln Leu Thr  
 65 70 75 80  
 45 Pro Leu Asn Leu Val Ser Ser Ala Asp Ala Thr His Gln Phe Leu Gly  
 85 90 95  
 Gly Glu Gly Gln Trp Leu Leu Ala Met Pro Ala His His Ile Ala Gly  
 100 105 110  
 50 Met Gln Val Leu Leu Arg Ser Leu Ile Ala Gly Val Glu Pro Leu Ala  
 115 120 125  
 55 Ile Asp Leu Ser Thr Gly Phe His Ile Asp Ala Phe Ala Gly Ala Ala  
 130 135 140  
 Ala Glu Leu Lys Asn Thr Gly Asp Arg Val Tyr Thr Ser Leu Thr Pro  
 145 150 155 160

000472-022304

Met Gln Leu Leu Lys Ala Met Asp Ser Leu Gln Gly Ile Glu Ala Leu  
165 170 175

5 Lys Leu Phe Asp Val Ile Leu Val Gly Gly Ala Ala Leu Ser Lys Gln  
180 185 190

Ala Arg Ile Ser Ala Glu Gln Leu Asp Ile Asn Ile Val Thr Thr Tyr  
195 200 205

10 Gly Ser Ser Glu Thr Ser Gly Gly Cys Val Tyr Asp Gly Lys Pro Ile  
210 215 220

Pro Gly Ala Lys Val Arg Ile Ser Asp Glu Arg Ile Glu Leu Gly Gly  
225 230 235 240

15 Pro Met Ile Ala Gln Gly Tyr Arg Asn Ala Pro Glu His Pro Asp Phe  
245 250 255

20 Ala Asn Glu Gly Trp Phe Thr Thr Ser Asp Ser Gly Glu Leu His Asp  
260 265 270

Gly Ile Leu Thr Val Thr Gly Arg Val Asp Thr Val Ile Asp Ser Gly  
275 280 285

25 Gly Leu Lys Leu His Pro Glu Val Leu Glu Arg Ala Ile Ala Asp Ile  
290 295 300

Lys Gly Val Thr Ala Ala Cys Val Val Gly Ile Pro Asp Pro Arg Leu  
305 310 315 320

30 Gly Gln Ala Ile Val Ala Ala Tyr Ser Gly Ser Ile Ser Pro Ser Glu  
325 330 335

35 Val Ile Glu Gly Leu Asp Asp Leu Pro Arg Trp Gln Leu Pro Lys Arg  
340 345 350

Leu Lys His Leu Glu Ser Leu Pro Ser Ile Gly Pro Gly Lys Ala Asp  
355 360 365

40 Arg Arg Ala Ile Ala Lys Leu Phe  
370 375

45

<210> 3  
<211> 19  
<212> DNA  
<213> Corynebacterium glutamicum

50

<220>  
<223> Primer menE-int1

<400> 3

55 ctcaactccgt tgaatttgg

<210> 4  
<211> 19

&lt;212&gt; DNA

&lt;213&gt; Corynebacterium glutamicum

&lt;220&gt;

5 &lt;223&gt; Primer menE-int2

&lt;400&gt; 4

caggtgcatt tctgtagcc

19

10

000551 BT / IP